

Preprint PNPI-1947, January 1994

Critical Composite Superconformal String

V.A.Kudryavtsev

Petersburg Nuclear Physics Institute
Gatchina, St.Petersburg 188350, Russia

Abstract

In the paper the nilpotent conditions of BRST operator for new superconformal string model were found. This string includes anticommutation $2-d$ fields additional to the standard Neveu-Schwarz superconformal set which carry quark quantum numbers. In this case the superconformal symmetry is realized by a non-linear way. In the superconformal composite string new constraints for 1 and $1/2$ conformal dimension should be added to the standard system of Virasoro superalgebra constraints for 2 and $3/2$ conformal dimensions. The number N of the constraints and numbers D and D' of bosonic and fermionic $2-d$ fields are connected by a simple relationship $D'/2 + D - 3N - 15 = 0$. Also perspectives of the critical composite superconformal string are discussed.

Four dimensional string models [1] derived from a fermionization of compactification of $2 - d$ bosonic fields have essentially extended the field of critical string under consideration. The next step forward in this direction has been done in the recently proposed model of superfermionic superconformal string [2]. In the superfermionic string all the $2 - d$ bosonic fields are excluded and only fermionic ones are conserved. Such an approach revealed some interesting possibilities for real describing quark degrees of freedom in the case of hadrons, but it ceased to have a remarkable feature of duality of amplitude characteristic for all original string theories [3]. Synthesis of the classical approach and the superfermionic model [4] based upon the generalization of multi-Regge vertices [5] proved to be more promising. The generalization allowed us to build a realistic model of π -meson interactions possessed spectrum of resonance states, Regge trajectories and lengths of scattering in a good agreement with experiment. In the composite superconformal model the most attractive features of the original string approach (superconformal symmetry and the duality) are conserved.

In the present paper we find how one can obtain the composite superconformal model with critical properties. As known, the standard Neveu-Schwarz-Ramond string (NSR) is critical in 10-d space-time. The composite string in addition to the standard NSR fields [6] $X_m(\tau)$ and $H_m(\tau)$, $m = 0, 1, 2 \dots, d - 1$ (a string coordinate and its superpartner) has a supplementary set of 36 anticommutation fields of conformal dimension $\frac{1}{2}$ [2] in a superfermionic sector. They are 8-component fields $\Psi_{\alpha\beta}$ (α — a spinor index and β — isospinor one, $\alpha = 1, 2, 3, 4$; $\beta = 1, 2$) carrying quantum numbers of an isospinor quark with the spin $\frac{1}{2}$ and also 28 fields with quantum numbers of currents built from these quarks. Among the 28 fields we have 10 isoscalar fields (4 vectors φ_μ , 6 tensors $\xi_{\mu\nu}$) and 18 isovector fields (3 scalars η_i , 3 pseudoscalars v_i and 12 pseudovectors $\chi_{\mu i}$, $\mu = 0, 1, 2, 3$; $i = 1, 2$). These 36 components appear in the three-linear (in these fields) scalar supergauge operator G_r^f of Virasoro algebra ($G_r^f \equiv G_r^{(0)}$ in notations of ref.[2]). The sum of G_r^f and the standard Neveu-Schwarz operator $G_r^{NS} = \oint \partial\tau X_\mu H^\mu e^{-in\tau}$ gives a total supergauge operator G_r of the considered model

$$G_r = G_r^f + G_r^{NS} \quad (1)$$

which satisfies the anticommutation relations of the standard Virasoro su-

peralgebra

$$\begin{aligned}
\{G_r, G_s\} &= 2L_{r+s} + \frac{c}{3} \left(r^2 - \frac{1}{4}\right) \delta_{r,-s} \\
[L_n, L_m] &= (n-m) L_{n+m} + \frac{c}{12} n(n^2-1) \delta_{n,-m} \\
[L_n, G_r] &= \left(\frac{n}{2} - r\right) G_{n+r}
\end{aligned} \tag{2}$$

where

$$c = \frac{3d}{2} + 18 \quad . \tag{3}$$

Wave vectors of physical states $|\Phi\rangle$ satisfy the corresponding gauge conditions

$$\begin{aligned}
G_r |\Phi\rangle &= 0 & r > 0 \\
L_n |\Phi\rangle &= 0 & n > 0 \\
(L_0 - a) |\Phi\rangle &= 0
\end{aligned} \tag{4}$$

where a constant $a = \frac{1}{2}$ in the NSR model.

For the standard NS superstring the critical value of c , which provides unitarity in loop corrections, is equal to 15 (that is $d = 10$ in the absence of the superfermionic sector). For $d \geq 0$ in the composite superconformal model with the conditions (4) the value (3) is not critical and, consequently, the unitarity on loop level is lost. However, as we shall see, one can achieve the critical supergauge algebra of the constraints by having involved additional constraints

$$\begin{aligned}
\nu_r^{(l)} |\Phi\rangle &= 0 & r > 0 \\
J_n^{(l)} |\Phi\rangle &= 0 & n > 0
\end{aligned} \quad l = 1, 2, \dots, N \tag{5}$$

where $\nu_r^{(l)}$ some N linear combinations of fields from the considered above 36 components of the superfermionic sector and $J_n^{(l)} = \{G_{n-r}, \nu_r^{(l)}\}$ are their associated currents. Certainly, it also means the requirement of invariance of string vertices under the new constraints (5). The incorporation of the new gauge constraints (5) produces new items which depend on new ghost and antighost operators $d_n^{(l)}, \ell_n^{(l)}, \delta_r^{(l)}$ and $\alpha_r^{(l)}$ in BRST charge Q according to

commutation relations for $\nu_r^{(l)}$ and $J_n^{(l)}$

$$\begin{aligned}
[J_n^{(i)}, J_m^{(j)}] &= f_{ijk} J_{n+m}^{(k)} + g_{ij} n \delta_{n,-m} \\
[L_n, J_m^{(i)}] &= -m J_{n+m}^{(i)} \\
\{G_r, \nu_s^{(i)}\} &= J_{r+s}^{(i)} \\
[L_n, \nu_s^{(i)}] &= \left(-\frac{n}{2} - s\right) \nu_{n+s}^{(i)} \\
[J_n^{(i)}, \nu_s^{(j)}] &= t_{ijl} \nu_{n+s}^{(l)} \\
\{\nu_s^{(i)}, \nu_r^{(j)}\} &= h_{ij} \delta_{r,-s}
\end{aligned} \tag{6}$$

If the corresponding operator of the BRST charge Q for the composite superconformal string is written in full, then we have

$$\begin{aligned}
Q &= \sum_n L_{-n} c_n + \sum_r G_{-r} \gamma_r + \sum_n J_{-n} d_n^i + \sum_r \nu_{-r}^{(j)} \delta_r^{(j)} \\
&- \frac{1}{2} \sum_{n,m} (m-n) : c_{-m} c_{-n} b_{m+n} : - \sum_{n,r} \left(\frac{3n}{2} + r\right) : c_{-n} \beta_{-r} \gamma_{r+n} : \\
&- \sum_{r,s} \gamma_{-r} \gamma_{-s} b_{r+s} - \frac{1}{2} \sum_{i,j,k,n,m} f_{ijk} : d_{-n}^{(i)} d_{-m}^{(j)} \ell_{n+m}^{(k)} : \\
&+ \sum_{n,m} (n+m) c_{-n} \ell_{-m}^{(i)} d_{m+n}^{(i)} - \sum_{r,m} \gamma_{-r} \ell_{-m}^{(i)} \delta_{r+m}^{(i)} \\
&- \sum_{i,r,s} (r+s) \gamma_{-r} \alpha_{-s}^{(i)} d_{r+s}^{(i)} - \sum_{n,s} \left(\frac{n}{2} + s\right) c_{-n} \alpha_{-s}^{(i)} \delta_{n+s}^{(i)} \\
&- \sum_{i,l,j} t_{ijl} d_{-n}^{(i)} \alpha_{-s}^{(j)} \delta_{n+s}^{(l)} + \sum_i f^{(i)} d_0^{(i)} - a c_0
\end{aligned} \tag{7}$$

where

$$\begin{aligned}
\{c_n, b_m\} &= \delta_{n,-m} & [\beta_r, \gamma_s] &= \delta_{r,-s} \\
c_n^+ &= c_n & b_m^+ &= b_{-m} & \ell_n^{(i)+} &= \ell_{-n}^{(i)} & d_n^{(i)+} &= d_{-n}^{(i)} \\
\beta_r^+ &= -\beta_{-r} & \gamma_r^+ &= \gamma_{-r} & \alpha_r^{(i)+} &= -\alpha_{-r}^{(i)} & \delta_s^{(j)+} &= \delta_{-s}^{(j)}
\end{aligned}$$

$$\left[\alpha_r^{(i)}, \delta_s^{(j)}\right] = \delta_{i,j} \delta_{r,-s} \quad \left\{\ell_n^{(i)}, d_m^{(j)}\right\} = \delta_{i,j} \delta_{n,-m}$$

Now, as for BRST approach in the original string theories one should find conditions that the BRST charge Q is nilpotent, which means that the superalgebra of the gauge constraints is critical one. As a result of this nilpotency $Q^2 = 0$ anomal terms are absent in the commutation relations (8) and (9).

$$\left[L_n^{quant}, L_m^{quant}\right] = (n - m) L_{n+m}^{quant} \quad (8)$$

$$\left\{G_r^{quant}, G_s^{quant}\right\} = 2L_{r+s}^{quant} \quad (9)$$

where $L_n^{quant} = \{b_n, Q\}$ and $G_s^{quant} = [\beta_s, Q]$

From disappearance of anomalies in (8) and (9) we obtain two relationships

$$D + \frac{D'}{2} - 3N - 15 = 0 \quad (10)$$

$$a = \frac{1}{2} \quad (11)$$

where D is a number of bosonic fields and D' is a number of fermionic fields. In this case we have $D = d$, and $D' = d + 36$.

The relationship (10) becomes readily available when we know contributions of ghosts of conformal dimension j in the anomaly (8): $(\pm 1) (3(2j - 1)^2 - 1)$. The sign $\pm$ is defined by statistics of the ghost. We have $(-1) 26$ for $j = 2$, $(+1) (+11)$ for $j = \frac{3}{2}$, $(-1) (+2)$ for $j = 1$, and $(+1) (-1)$ for $j = \frac{1}{2}$ that allows us to obtain

$$-26 + 11 - 2N - N = -15 - 3N .$$

A concrete method of construction of string vertices and amplitude of the composite superconformal string [4] implies that both "gluonic" Neveu-Schwarz sector and "quark" (superfermionic) sector are individually critical. So we choose here $d = 10$ and $N = 6$. Furthermore, it appears that the conditions of absence of anomalies don't insure $Q^2 = 0$ because of anomalies in the commutation relations $\left\{\nu_s^{(i)}, \nu_r^{(j)}\right\}$ and $\left[J_n^{(i)}, J_m^{(j)}\right]$ in (6). One can check that the nilpotent condition $Q^2 = 0$ requires additional constraints

$$\begin{aligned} f^{(i)} &= 0 & h_{ij} &= 0 & \sum_j f_{ijj} &= 0 \\ \sum_j t_{jij} &= 0 & \sum_j t_{ijj} &= 0 \end{aligned} \quad (12)$$

The disappearance of coefficients h_{ij} is possible due to a signature of Minkowski metric and existence of the components $\varphi_0, \xi_{0\mu}, \chi_{0i}$ in the spectrum of the superfermionic sector

$$\begin{aligned} \{(\varphi_\mu)_r, (\varphi_\nu)_s\} &= -g_{\mu\nu}\delta_{r,-s} \\ g_{\mu\nu} &= \text{diag}(1, -1, -1, -1) \end{aligned} \quad (13)$$

Choose, as $\nu^{(l)}$ $l = 1, 2, \dots, N$, six light-like components $\varphi_0 + \varphi_3, \xi_{01} + \xi_{31}, \xi_{02} + \xi_{32}, \chi_{0i} + \chi_{3i}$ ($i = 1, 2, 3$). By choosing so one can obtain $Q^2 = 0$ that is the desired condition. It is remarkable that in this case, as for the original critical string (the bosonic string ($d = 26$) and the superstring ($d = 10$)), an elimination of spurious states with null norm has occurred together with their conjugate ones. This leads to vanishing both null and longitudinal components. In other words the elimination of $\varphi_0 + \varphi_3, \xi_{01} + \xi_{31}, \xi_{02} + \xi_{32}, \chi_{0i} + \chi_{3i}$ occurs together with their conjugate $\varphi_0 - \varphi_3, \xi_{01} - \xi_{31}, \xi_{02} - \xi_{32}, \chi_{0i} - \chi_{3i}$. Hence, $\varphi_0, \varphi_3, \xi_{01}, \xi_{31}, \xi_{02}, \xi_{32}, \chi_{0i}, \chi_{3i}$ fall out from spectrum of physical states. Moreover, in the critical composite superconformal model the elimination of spurious states appears to be still more effective by virtue of the nontrivial commutation relations (6) for these components and currents connected with them. It is interesting that in the string amplitude [4] one can choose as light-like ($k_i^2 = 0$) all momenta along the quark lines of dual quark diagram[7]. Since the superfermionic sector in the amplitudes [4] is chosen for each quark lines, this is in its turn achieving a necessary light direction for the considered above conditions $\varphi_0 + \varphi_3 = k_\mu \varphi^\mu$ and so on, and hence we insure Lorentz invariance of this approach.

The analysis which has been out here corresponds to the first generation (u and d quarks of the Standard Model). Other quantum numbers which correspond to s, c, b and t quarks and baryon number will be a subject of a following paper. It is to be noted that the number of the fermionic fields exceeds the number of bosonic ones in the spectrum of the composite superconformal string, and this fact together with absence of tachions and dilatons inevitably has to lead to finiteness of oriented (only possible here) one-loop (and, perhaps, multi-loop) diagrams. As we know the leading divergence of integral in the Chan variable x for the one-loop diagram is defined by the expression $\exp\left(\frac{\pi^2}{6}(\widetilde{D} - \widetilde{D}')\right) \frac{1}{1-x}$ in the limit $x \rightarrow 1$ where $\widetilde{D}, \widetilde{D}'$ -full numbers of boson and fermion fields correspondingly including ghost fields. In the case of $\widetilde{D} < \widetilde{D}'$ and $x \leq 1$ this integral converges.

The author would like to thank E.N.Antonov, A.P.Bukhvostov, G.V.Frolov, L.N.Lipatov for useful discussions.

Work supported in part by Russian Fund of Fundamental Research under Contract No.93-02-3864

References

- [1] I.Antoniadis, C.P.Bachas, C.Kounnas, *Nucl. Phys.*, **B289**, 87 (1987)
- [2] V.A.Kudryavtsev, *Pis'ma Zh. Eksp. Teor. Fiz.*, **54**, 417 (1991)
V.A.Kudryavtsev, *Yad. Fiz.*, to be published.
- [3] M.B.Green, J.H.Schwarz and E.Witten, *Superstring Theory* (Cambridge Univ. Press, N.Y. 1987)
- [4] V.A.Kudryavtsev, *Pis'ma Zh. Eksp. Teor. Fiz.*, **58**, 321 (1993)
- [5] C.Lovelace, *Phys. Lett.*, **B32**, 490 (1970)
D.Olive, *Nuovo cim.*, **A3**, 399 (1971)
- [6] A.Neveu, J.H.Schwarz,C.B.Thorn, *Phys. Lett.*, **B35**, 529 (1971)
- [7] H.Harari, *Phys. Rev. Lett.*, **22**, 562 (1969)